

<b>REPORT DOCUMENTATION PAGE</b>				Form Approved OMB No. 0704-0188		
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Service, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188) Washington, DC 20503.						
<b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</b>						
<b>1. REPORT DATE (DD-MM-YYYY)</b> 15-10-2002		<b>2. REPORT DATE</b> FINAL grant report		<b>3. DATES COVERED (From - To)</b> 18-08-1998 to 31-01-2002		
<b>4. TITLE AND SUBTITLE</b> An Innovative Coastal-Ocean Observing Network (ICON)				<b>5a. CONTRACT NUMBER</b>		
				<b>5b. GRANT NUMBER</b> <b>N00014-98-1-0870</b>		
				<b>5c. PROGRAM ELEMENT NUMBER</b>		
<b>6. AUTHOR(S)</b> Fernandez, Daniel M.; Paduan, Jeffrey D.; Chavez, Francisco P; Shulman, Igor; Vesecky, John F.; Kindie, John C.; Maffione, Robert; Barrick, Donald E.				<b>5d. PROJECT NUMBER</b>		
				<b>5e. TASK NUMBER</b>		
				<b>5f. WORK UNIT NUMBER</b>		
<b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b> California State University, Monterey Bay 100 Campus Center Seaside, CA 93955				<b>8. PERFORMING ORGANIZATION REPORT NUMBER</b>  F9910040		
<b>9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)</b> Foundation of California State University, Monterey Bay 100 Campus Center, Bldg. 86C Seaside, CA 93955				<b>10. SPONSOR/MONITOR'S ACRONYM(S)</b>  CSUMB/FCSUMB		
				<b>11. SPONSORING/MONITORING AGENCY REPORT NUMBER</b>		
<div style="display: flex; justify-content: space-between;"> <div> <b>12. DISTRIBUTION AVAILABILITY STATEMENT</b>            No restrictions necessary         </div> <div style="text-align: center;"> <b>DISTRIBUTION STATEMENT A</b>            Approved for Public Release            Distribution Unlimited         </div> <div style="font-size: 2em; font-weight: bold;">           20021023 077         </div> </div>						
<b>13. SUPPLEMENTARY NOTES</b>						
<b>14. ABSTRACT</b> The Innovative Coastal-Ocean Observing Network (ICON) is a partnership of government, academic, and industrial entities funded by the National Ocean Partnership Program (NOPP). Its goal is to bring together modern measurement technologies, to develop new technologies, and to integrate them within a data assimilating coastal ocean circulation model. The major components of the observing network include 1) surface current maps from shore-based high frequency (HF) radar installations, 2) subsurface currents, temperature, salinity, and bio-optical properties plus surface meteorological properties from several deep-ocean moorings, 3) sea surface temperature and color from satellites, and 4) along-track temperature and temperature variances from two acoustic tomography slices through the region. The model performance is quite good at seasonal time scales, which is a validation of the one-way nesting because these variations are successfully tracked by the PWC regional-scale model. At higher frequencies, the model does not reproduce the observed level of variability. For the alongshore currents down to 150 meter depth, data assimilation resulted in greater correlation between modeled and observed currents.						
<b>15. SUBJECT TERMS</b> <div style="display: flex; justify-content: space-around; margin-top: 10px;"> <div>data assimilation coastal ocean circulation</div> <div>ocean modeling high-frequency radar</div> <div>oceanographic data collection bio-optical sensors</div> <div>acoustic tomography</div> </div>						
<b>16. SECURITY CLASSIFICATION OF:</b>			<b>17. LIMITATION OF ABSTRACT</b>		<b>18. NUMBER OF PAGES</b>	
a. REPORT  UU	b. ABSTRACT  UU	c. THIS PAGE  UU	UU		7 pages follow	
<b>19a. NAME OF RESPONSIBLE PERSON</b> Daniel M. Fernandez/Cynthia E. Lopez						
<b>19b. TELEPHONE NUMBER (Include area code)</b> (831) 582-3786/(831) 582-3089						

**Grant: N00014-98-1-0870**

**Grantee: Foundation of California State University, Monterey Bay**

**Title: "An Innovative Coastal-Ocean Observing Network (ICON)"**

(The Naval Postgraduate School was also a grantee under this grant and also submitted a similar report)

Jeffrey D. Paduan

Steven R. Ramp, Leslie K. Rosenfeld, Curtis A. Collins, Ching-Sang Chiu, Newell Garfield  
Department of Oceanography, Code OC/Pd, Naval Postgraduate School

Monterey, CA 93943

phone: (831) 656-3350; fax: (831) 656-2712; email: paduan@nps.navy.mil

Daniel M. Fernandez

Institute of Earth Systems Science & Policy, California State University Monterey Bay  
Seaside, CA 93955

phone: (831) 582-3786; fax: 582-4688; email: daniel\_fernandez@csumb.edu

Francisco P. Chavez

Monterey Bay Aquarium Research Institute

7700 Sandholdt Road

Moss Landing, CA 95039

phone: (831) 775-1709; fax: (831) 775-1620; email: chfr@mbari.org

Igor Shulman

Institute of Marine Sciences, University of Southern Mississippi

Stennis Space Center, MS 39529

phone: (228) 688-3403; fax: (228) 688-7072; email: shulman@coam.usm.edu

John F. Vesecky

Atmospheric, Oceanic, & Space Sciences, University of Michigan

Ann Arbor, MI 48109

phone: (734) 764-5151; fax: (734) 764-5137; email: jfv@engin.umich.edu

John C. Kindle

Oceanography Division, Naval Research Laboratory

Stennis Space Center, MS 39529

phone: (228) 688-4118; fax: (228) 688-4759; email: kindle@nrlssc.navy.mil

Robert Maffione

HOBILabs

56 Penny Lane, Suite 104, Watsonville, CA 95076

(831) 768-0680; fax: (831) 768-0681; email: hobilabs@hobilabs.com

Donald Barrick

1000 Freemont Ave, Suite K, Los Altos, CA 94024

phone: (408) 773-8240; fax: (408) 773-0514; email: don@codaros.com

## **LONG-TERM GOALS**

The Innovative Coastal-Ocean Observing Network (ICON) is a partnership of government, academic, and industrial entities funded by the National Ocean Partnership Program (NOPP). Its goal is to bring together modern measurement technologies, to develop new technologies, and to integrate them within a data assimilating coastal ocean circulation model.

## **OBJECTIVES**

The objectives of the project are to evaluate the several real-time observing systems as components of future coastal monitoring networks as well as sources for data-assimilating numerical models.

## **APPROACH**


The approach taken in this project is to build on existing partnerships and observing systems around the Monterey Bay region by providing coordination, additional instrumentation, and a focus on evaluating the impact of the various measurements on the validation and forcing of a coastal circulation model. The major components of the observing network include 1) surface current maps from shore-based high frequency (HF) radar installations, 2) subsurface currents, temperature, salinity, and bio-optical properties plus surface meteorological properties from several deep-ocean moorings, 3) sea surface temperature and color from satellites, and 4) along-track temperature and temperature variances from two acoustic tomography slices through the region. These data sets each involve real-time data telemetry. They are also being used as either validation or assimilation sources for a nested, primitive equation numerical model designed to track the evolution of mesoscale filaments and eddies.

## **WORK COMPLETED**

Additional HF radar stations were deployed south of Monterey Bay, moored observations were retrieved from five offshore sites, and acoustic travel times were collected from one along-shore and one cross-shore tomography section. The mooring locations, HF radar sites, and a sample HF radar data set are shown in Figure 1. Of particular note are expanded HF radar locations at Granite Canyon and Pt. Sur and the M4 mooring location, which returned exceptionally good oceanographic, atmospheric, and bio-optical data for the entire period of its deployment from August 1999 into September 2000. Shorter-term deployments of the NPS FLUX buoy and of a bottom-mounted ADCP with acoustic telemetry were made in August 2000 in conjunction with the ONR/AOSN field campaign. Also during that campaign, thirteen separate synoptic surveys of sea surface temperature, air temperature, and winds were made from an instrumented aircraft.

The numerical model was spun up from 1995 using NOGAPS wind forcing and lateral boundary conditions from the NRL Pacific West Coast (PWC) model, which itself is nested within the NRL 1/4 degree, global layered model. Multiple runs of the ICON model for the period 1999 have been conducted to evaluate the role of wind forcing (NOGAPS vs. COAMPS), the effectiveness of various data assimilation techniques, and the sensitivity to surface heat fluxes. In addition, a triply nested

model grid was generated for the Monterey Bay region to be used to investigate specific case studies of upwelling fronts and filaments sampled during the AOSN field campaign.




*Figure 1. ICON Mooring and radar locations presented with the first five-site HF radar data set collected on 20 January 2000. Note that these single-component, radial currents must still be combined to produce a vector current map in the regions of overlap.*

## RESULTS

The HF radar data in Figure 1 illustrate the results of the expanded HF radar network. Data from the entire network is available beginning January 2000. Analyses to date have, however, utilized surface current maps from the three sites around Monterey Bay. Model results, with and without assimilation of the HF radar-derived currents, have been compared to temperatures, salinities, and velocities at the mooring sites. In addition, a systematic comparison has been initiated to evaluate the surface heat flux parameters output by the high-resolution (9 km) COAMPS atmospheric model against time series measurements from the mooring sites.

As an example of the model performance and types of comparisons that have been conducted, Figure 2 shows subsurface temperatures for all of 1999 at the M1 mooring location from observations and from

the numerical model. Statistical comparisons of the same data are presented in Table 1 for the base run with NOGAPS wind forcing and no heat flux forcing and for a run in which surface heat fluxes were incorporated through the assimilation of the composite MCSST data.




**Figure 2.** Daily averaged temperatures at the M1 mooring site from observations and from the ICON model forced with NOGAPS winds and regional-scale (PWC) boundary conditions.

The model performance is quite good at seasonal time scales, which is a validation of the one-way nesting because these variations are successfully tracked by the PWC regional-scale model. At higher frequencies, the model does not reproduce the observed level of variability. Other problems are visible near the surface where the model mixed layers are too deep in summer and too shallow in winter, which is certainly a reflection of the lack of heat flux forcing. This problem is mitigated to some degree by assimilation of MCSST data at the surface (Table 1).

**Table 1.** Model versus observed temperature statistics ( $^{\circ}\text{C}$ ) at the M1 mooring site for 1999.

Depth (m)	$\sigma_{\text{obs}}$	$\sigma_{\text{mod}}$	$\sigma_{\text{mod}}$ with MCSST	RMS difference	RMS diff. with MCSST
0	1.52	1.12	1.16	1.33	1.05
20	1.03	1.12	1.07	1.85	1.72
60	0.81	0.83	0.83	1.59	1.58
100	0.70	0.69	0.66	1.28	1.23
200	0.46	0.47	0.46	0.72	0.63
300	0.30	0.35	0.35	0.38	0.35


Assimilation of velocity data from low-pass-filtered HF radar data has been tested using the machinery of Physical-Space Statistical Analysis System (PSAS) and spatial error covariances based on seasonal statistics from the HF radar data (see Shulman et al., 2000ab). Despite the surface-only nature of the assimilation (no vertical projection was made during the assimilation), divergences in the correction fields imposed by the observed velocity patterns quickly penetrate to the main thermocline. This is illustrated in Figure 3, which gives an example of the model currents at 75m with and without data assimilation. The changes at 75m are significant and, in addition, the limited-area corrections can be seen to have an effect outside the spatial range of the actual data.



**Figure 3.** *ICON model results at 75m depth on 15 August 1995 with only (NOGAPS) wind forcing (a) and with the addition of HF radar-derived surface current assimilation (b). The model difference at 75 m depth with and without assimilation (c) and the most recent HF radar-derived surface current map (d) are also shown.*

Long-term velocity comparisons at the M1 mooring location illustrate the effect of surface velocity data assimilation as a function of depth. Correlations of observed and modeled cross-shore (u) and along-shore (v) velocities are shown in Figure 4 as a function of depth with and without data assimilation. The figure shows largely improved results down to 150m for the dominant along-shore

currents when data assimilation is employed. The results are not all good as the correlation with weaker cross-shore currents is reduced in the region of the thermocline with data assimilation.



*Figure 4. Modeled vs. observed currents at M1 in 1995.*

## IMPACT/APPLICATIONS

The likely impacts of this project include improved real-time communication, processing, and display of coastal ocean data along with improved algorithms for assimilating that data into numerical models.

## TRANSITIONS

The transition opportunities are related to improved coastal nowcast and forecast systems.

## RELATED PROJECTS

This project is closely related to other NOPP efforts focusing on data assimilation and coastal ocean modeling. The ONR-sponsored project of Ly and Paduan (N00014-00-WR20098) to observe and model surface waves in Monterey Bay is a direct extension of the ICON efforts. Also, related efforts can be found in the separate modeling program of Shulman (N00014-97-1-0171). ICON is closely linked to the Autonomous Ocean Sampling Network (AOSN) project whose field program, along with a campaign sponsored by the Monterey Bay Aquarium Research Institute called MUSE, was conducted in Monterey Bay in August 2000.



## PUBLICATIONS

Barrick, D., R. Cheng, N. Garfield, J. Paduan, P. Lilleboe, J. Gartner, and L. Pederson, 2000: Toward bay/harbor circulation model improvement incorporating HF radar data based on SeaSonde deployments on San Francisco Bay. Proceedings, IEEE Oceans-2000, 5 pp.

Paduan, J.D., L.K. Rosenfeld, S.R. Ramp, F. Chavez, C.S. Chiu, and C.A. Collins, 1999: Development and maintenance of the ICON observing system in Monterey Bay. Proceedings, Third Conference on Coastal Atmospheric and Oceanic Prediction and Processes, New Orleans, LA, 3-5 November.

Shulman, I., C.-R. Wu, J.K. Lewis, J.D. Paduan, L.K. Rosenfeld, S.R. Ramp, M.S. Cook, J.C. Kindle, and D.-S. Ko, 2000a: Development of the high resolution, data assimilating numerical model of the Monterey Bay. In Spaulding, M.L. and H. Lee Butler (Ed.), *Estuarine and Coastal Modeling*, 980-994.

Shulman, I., C.-R. Wu, J.K. Lewis, J.D. Paduan, L.K. Rosenfeld, J.C. Kindle, S.R. Ramp, and C.A. Collins, 2000b: High resolution modeling and data assimilation in the Monterey Bay area. Continental Shelf Research, submitted.